



ELSEVIER

Atherosclerosis 170 (2003) 13–19

ATHEROSCLEROSIS

www.elsevier.com/locate/atherosclerosis

## A casein variant in cow's milk is atherogenic

Kristy A. Tailford, Celia L. Berry, Anita C. Thomas, Julie H. Campbell\*

Centre for Research in Vascular Biology, School of Biomedical Sciences, University of Queensland, Brisbane, Qld 4072, Australia

Received 9 November 2002; received in revised form 25 February 2003; accepted 31 March 2003

### Abstract

Casein is a major protein in cow's milk that occurs in several variant forms, two of which are  $\beta$ -casein A<sup>1</sup> and  $\beta$ -casein A<sup>2</sup>. The levels of these two proteins vary considerably in milk dependent on the breed of cow, and epidemiology studies suggest that there is a relationship between their consumption and the degree of atherosclerosis. In the present study, the direct effect of consumption of  $\beta$ -casein A<sup>1</sup> vs  $\beta$ -casein A<sup>2</sup> on atherosclerosis development was examined in a rabbit model. Sixty rabbits had their right carotid artery balloon de-endothelialised at  $t = 0$ , divided randomly into 10 groups ( $n = 6$  per group), then for 6 weeks fed a diet containing 0, 5, 10 or 20% casein isolate, either  $\beta$ -casein variant A<sup>1</sup> or A<sup>2</sup>, made up to 20% milk protein with whey. Some groups had their diets supplemented with 0.5% cholesterol. Blood samples were collected at  $t = 0, 3$  and 6 weeks and rabbits were sacrificed at  $t = 6$  weeks. In the absence of dietary cholesterol,  $\beta$ -casein A<sup>1</sup> produced significantly higher ( $P < 0.05$ ) serum cholesterol, LDL, HDL and triglyceride levels than whey diet alone, which in turn produced higher levels than  $\beta$ -casein A<sup>2</sup>. Rabbits fed  $\beta$ -casein A<sup>1</sup> had a higher percent surface area of aorta covered by fatty streaks than those fed  $\beta$ -casein A<sup>2</sup> ( $5.2 \pm 0.81$  vs  $1.1 \pm 0.39$ ,  $P < 0.05$ ) and the thickness of the fatty streak lesions in the aortic arch was significantly higher ( $0.04 \pm 0.010$  vs  $0.00$ ,  $P < 0.05$ ). Similarly, the intima to media ratio (I:M) of the balloon injured carotid arteries in A<sup>1</sup> fed animals ( $0.77 \pm 0.07$ ) was higher than in those that consumed A<sup>2</sup> ( $0.57 \pm 0.04$ ) or whey ( $0.58 \pm 0.04$ ), but this did not reach significance. In the presence of 0.5% dietary cholesterol, the thickness of the aortic arch lesions was higher ( $P < 0.05$ ) in 5, 10 and 20% casein A<sup>1</sup> fed animals compared with their A<sup>2</sup> counterparts, while other parameters were not significantly different. It is concluded that  $\beta$ -casein A<sup>1</sup> is atherogenic compared with  $\beta$ -casein A<sup>2</sup>.

© 2003 Elsevier Science Ireland Ltd. All rights reserved.

Keywords:  $\beta$ -Casein A<sup>1</sup>;  $\beta$ -Casein A<sup>2</sup>; Atherosclerosis; Cow milk; Fatty streaks

### 1. Introduction

Cow milk constitutes a major part of the diet of developed nations and it contains an average 32 g of protein per litre [1], mostly casein (82%) and whey (18%). The caseins have been divided into four subclasses,  $\alpha_{s1}$ ,  $\alpha_{s2}$ ,  $\beta$  and  $\kappa$  [2,3], whose major function is to transport calcium phosphate [4].  $\beta$ -Casein is the second most abundant casein behind  $\alpha_{s1}$  [1] and is essential for curd formation and important in determining the surface properties of micelles [4]. To date 10 genetic variants of  $\beta$ -casein have been characterised with regard to sequence [5,6], and of these only A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and B are found in nearly all *Bos taurus* populations [7,8]. The allelic distribution of variants is such that the majority

of cows in Western nations produce milk rich in the genetic variant  $\beta$ -casein A<sup>1</sup>. For example, Holstein and Ayrshire breeds of cow produce 63 and 67% A<sup>1</sup>, respectively, and 35 and 33% A<sup>2</sup> [9]. In contrast, the Guernsey breed produces 1% A<sup>1</sup> and 98% A<sup>2</sup>. Alignment of  $\beta$ -casein A<sup>1</sup> and  $\beta$ -casein A<sup>2</sup> protein sequences reveals near-perfect homology with the only difference at codon 67 where  $\beta$ -casein A<sup>2</sup> has a proline residue and  $\beta$ -casein A<sup>1</sup> has a histidine residue [4,10].

While some epidemiological studies show no association between reported milk consumption and cardiovascular mortality [11], comparative data from several populations have suggested a strong correlation between the consumption of  $\beta$ -casein A<sup>1</sup> and deaths from ischaemic heart disease. McLachlan [12] examined the calculated consumption of  $\beta$ -casein A<sup>1</sup> (excluding cheese) against IHD mortality in males aged 30–69 in 16 developed nations and found a correlation value ( $r^2$ ) of 0.86. Upon comparing IHD and  $\beta$ -casein A<sup>1</sup> con-

\* Corresponding author. Tel.: +61-7-33654658; fax: +61-7-33651299.

E-mail address: julie.campbell@uq.edu.au (J.H. Campbell).

sumption in the states of the former West Germany, the same researcher found an  $r^2 = 0.66$ , three times greater than the correlation between smoking and IHD in the same population. This implies that  $\beta$ -casein A<sup>1</sup> consumption may cause more vascular damage than a well-known risk factor for atherosclerosis.

There have been almost no cases of IHD-related mortality reported in either the Masai (East Africa) [13] or Samburu (Northern Kenya) [14] communities. Furthermore, these tribes showed normal to low cholesterol levels despite having a diet rich in animal milk and meat protein [13,14]. Interestingly, almost all of the African Zebu (*B. indicus*) cattle populations produce milk containing  $\beta$ -casein A<sup>2</sup> [15].

Ingestion of total casein, the protein fraction isolated by acid precipitation of all casein subclasses from milk, has been shown to promote atherosclerosis in several animal models including rabbits [16], monkeys [17] and mice [18]. Casein consumption is also associated with hypercholesterolemia [19] and reduced LDL catabolism [20]. However, to date no controlled diet studies have been carried out to definitively determine whether a specific casein subclass or subclass variant is more atherogenic than others. The aim of the present study was to determine whether dietary administration of  $\beta$ -casein A<sup>1</sup> in a rabbit model of atherosclerosis promotes the disease state compared with rabbits fed  $\beta$ -casein A<sup>2</sup>.

## 2. Materials and methods

### 2.1. Study groups

Sixty New Zealand white/Lop cross rabbits of both sexes (16–24 weeks old) had their right carotid artery balloon de-endothelialised at  $t = 0$  [21], then randomly divided into 10 groups ( $n = 6$  per group). Group 1 was fed pellets containing 20% whey protein for 6 weeks while Groups 2 and 3 had 10%  $\beta$ -casein A<sup>1</sup> and A<sup>2</sup>, respectively, plus 10% whey, all with no added dietary cholesterol. The diets for the remaining seven groups contained 0.5% dietary cholesterol and 20% milk protein, which differed only in the proportions of whey and casein. Group 4 had 20% whey protein while Groups 5 and 6 received diets containing 10%  $\beta$ -casein A<sup>1</sup> and A<sup>2</sup>, respectively, plus 10% whey. Groups 7 and 8 had 5%  $\beta$ -casein A<sup>1</sup> and A<sup>2</sup>, respectively, plus 15% whey protein, while Groups 9 and 10 received 20%  $\beta$ -casein A<sup>1</sup> and A<sup>2</sup>, respectively. All diet pellets were specially prepared by Dyets, Inc., Philadelphia, PA.

In a pilot study, it was found that it took a few days for the rabbits to become accustomed to their new diets that differed from the normal pellets in colour, smell, texture and weight. Most obviously, the study pellets were white while normal pellets were brown. Thus, for 5 days prior to surgery, all rabbits were weaned onto their

respective diets by mixing increasing amounts of study pellets (25, 50, 75 and 100%) with normal rabbit pellets. At  $t = 0$ , only study pellets were offered for the following 6 weeks and water was available ad libitum. The amount of pellets eaten by each rabbit was determined each day. All rabbits were weighed weekly, commencing the day of surgery until the week of sacrifice. Blood was collected from each rabbit prior to surgery, at the commencement of diet and at  $t = 3$  and 6 weeks. The serum supernatant was extracted and analysed at Queensland Medical Laboratories, Brisbane, for total serum cholesterol, triglycerides, HDL, LDL and homocysteine.

At  $t = 6$  weeks, all animals were euthanased by an overdose of pentobarbitone (325 mg/ml Lethobarb, Virbac Australia Pvt. Ltd., Vic.), then immediately perfused fixed with 4% formaldehyde (Asia Pacific Specialty Chemicals Ltd., NSW, Australia) in phosphate-buffered saline at 100 mmHg. The entire aorta, from the iliac bifurcation to the aortic arch, was exposed and removed. The right (injured) and left (control) common carotid arteries were fully exposed and removed.

### 2.2. *En face* Oil-Red-O staining for fatty streak lesions

The thoracic and abdominal aorta minus the aortic arch (approximately 16 cm) was opened longitudinally and cut into four segments. Each segment was sewn (lumen side up) onto small pieces of clear plastic to ensure they remained flat. The segments were then stained *en face* with a saturated solution of the lipophilic dye Oil-Red-O.

Images of each aortic segment were scanned using Desk Scan II (version 2.31a, Hewlett-Packard, USA) and both the lesion and total area measured using the ImageJ imaging software (version 1.23y, National Institutes of Health, USA). The results were expressed as the percentage of luminal surface area covered by fatty streak lesions.

### 2.3. *Histological sections* intima to media ratio

Three segments of 3–4 mm length were cut from each aortic arch, dehydrated, embedded in plastic, sectioned and stained with Toluidine Blue. Similarly, four or five segments each of 3–4 mm in length were cut from the same regions spanning the full length of the left (control) and the right (ballooned) carotid arteries, respectively, and prepared in the same manner.

All morphometric analyses of the intima to media ratios (I:M) of the ballooned and control carotid arteries and the aortic arch were performed using the Mocha Image Analysis system (version 1.1, Jandel Scientific, CA). Measurements were taken to determine the size (in pixels) of both the intima and media layers in all

segments of each artery. The ratio of intima to media was calculated and recorded for each of the three aortic arch segments per rabbit, the four control carotid segments and the five ballooned carotid segments.

#### 2.4. Statistics

All statistical analyses were performed using the SigmaStat (Jandel Scientific, CA) statistical software package. Comparison of data from the morphometric analyses were carried out with one-way multiple ANOVAs, using the Kruskal–Wallis analysis of variance on ranks test. In all statistical analyses, a *P*-value of less than 0.05 was considered significant. All data have been reported as mean  $\pm$  standard error of mean.

### 3. Results

#### 3.1. Rabbit diets and weights

While the amount of the study pellets eaten each day varied between rabbits, on group averages the amounts consumed varied by only 12 g per day over the 6 weeks experimental period, and there was no significant difference ( $P < 0.05$ ) between any of the groups (mean =  $30.05 \pm 0.9$  g per day). The addition of 0.5% cholesterol to the diets of Groups 4–10 did not influence the amount eaten, or did the type or amount of casein. Almost all rabbits lost weight over the 6 weeks of study (mean  $-5.62\%$ ) with no significant difference between groups ( $P < 0.05$ ).

#### 3.2. Serum cholesterol and LDL/HDL ratio

At week 6, Group 3 ( $A^2$ ) serum cholesterol was significantly lower than all groups including Groups 1 (10% whey) and 2 ( $A^2$ ) (Fig. 1A). Interestingly, Group 2 ( $A^1$ ) was higher than Group 1, but this was not significant. Of note, the serum cholesterol level in Group 10 (20%  $A^2$  plus 0.5% dietary cholesterol) was approximately half that of Group 4 (20% whey plus 0.5% dietary cholesterol) and Group 9 (20%  $A^1$  plus 0.5% dietary cholesterol ( $12.1 \pm 1.9$  mmol/l vs  $24.7 \pm 5.7$  and  $22.4 \pm 6.8$  mmol/l, respectively) (see Fig. 1A and Table 1).

Group 3 ( $A^2$ ) LDL levels were significantly lower than Groups 1 and 2. All groups with added dietary cholesterol (Groups 4–10) were not significantly different from each other. These results, however, may have been swamped by the relatively high level of dietary cholesterol (0.5%), whereas a lower cholesterol load may have shown more striking differences. Serum HDL followed the same pattern as serum LDL. HDL in Group 2 ( $A^1$ ) was significantly higher than Group 3 ( $A^2$ ) but not significantly different from Group 1. The LDL




Fig. 1. (A) Total serum cholesterol levels (mmol/l) of rabbit Groups 1–6 at week 6. (B) Serum LDL to HDL ratio at week 6. Group 1: 20% whey diet; Group 2: 10%  $\beta$ -casein  $A^1$ +10% whey; Group 3: 10%  $\beta$ -casein  $A^2$ +10% whey; Group 4: 20% whey+0.5% cholesterol; Group 5: 10%  $\beta$ -casein  $A^1$ +0.5% cholesterol; and Group 6: 10%  $\beta$ -casein  $A^2$ +0.5% cholesterol. \* $P < 0.05$ .

to HDL ratio of Group 3 ( $A^2$ ) was significantly lower than all other groups including Group 1 (20% whey) and Group 2 (10%  $\beta$ -casein  $A^1$ ), while the LDL to HDL ratio of all groups with dietary cholesterol (Groups 4–10) were not significantly different from each other (see Fig. 1B). However, there was a trend for higher LDL to HDL ratios with increasing doses of  $\beta$ -casein  $A^1$  with dietary cholesterol ( $4.19 \pm 0.57$ ,  $4.51 \pm 0.34$  and  $4.99 \pm 0.67$ ) and lower levels with increasing doses of  $\beta$ -casein  $A^2$  with dietary cholesterol ( $5.63 \pm 1.75$ ,  $4.68 \pm 0.37$  and  $4.66 \pm 0.25$ ), but this was not significant (see Fig. 1B, Table 1).

#### 3.3. Triglycerides and homocysteine

By week 6, the serum triglyceride levels for most groups had fallen from those at week 0, and showed no significant difference between groups. Serum homocysteine levels changed only slightly from those at week 0, with no significant differences between the groups (data not shown).

Table 1

Data from rabbit Groups 7–10 at week 6 showing total serum cholesterol, serum LDL to HDL ratio, percent luminal surface area of aorta covered by lipid-filled lesions, thickness of lesions in aortic arch expressed as intima to media ratio (I:M) and thickness of lesions in injured carotid artery expressed as intima to media ratio (I:M)

Group	Chol (mmol/l)	LDL/HDL	Lesion (%)	Aorta I:M	Carotid I:M
7	18.6±3.7	4.19±0.57	10.6±1.54	0.04±0.01*	0.75±0.08
8	18.4±5.5	5.63±1.75	5.2±1.04	0.00±0.00*	0.70±0.08
9	22.4±6.8	4.99±0.67	6.4±1.57	0.05±0.01**	0.78±0.08
10	12.1±1.9	4.66±0.25	2.2±0.48	0.00±0.00**	0.65±0.05

Group 7: 5%  $\beta$ -casein A<sup>1</sup>+15% whey+0.5% cholesterol; Group 8: 5%  $\beta$ -casein A<sup>2</sup>+15% whey+0.5% cholesterol; Group 9: 20%  $\beta$ -casein A<sup>1</sup>+0.5% cholesterol; and Group 10: 20%  $\beta$ -casein A<sup>2</sup>+0.5% cholesterol.

\*  $P < 0.05$ .

\*\*  $P < 0.05$ .

### 3.4. Aortic fatty streaks

Group 1 rabbits fed 20% whey had 0% aortic luminal surface area covered by plaque (Fig. 2A). Group 3 rabbits (10%  $\beta$ -casein A<sup>2</sup> plus 10% whey) had only 1.1±0.4% surface area covered by plaque, and this was not significantly higher than Group 1. In contrast, the plaque surface area of Group 2 rabbits fed 10%  $\beta$ -casein A<sup>1</sup> plus 10% whey was 5.2±0.8%. All animals given a 0.5% cholesterol-enriched diet had values that were not significantly different from each other, irrespective of casein variant, although 20%  $\beta$ -casein A<sup>2</sup> with dietary cholesterol produced lesions covering only 2.2±0.48% of the aortic luminal surface compared with 6.4±1.57% for 20%  $\beta$ -casein A<sup>1</sup> under the same conditions (see Table 1). As with the other parameters measured, the addition of a dietary cholesterol load less than 0.5% may have produced more striking results. Interestingly, 10%  $\beta$ -casein A<sup>1</sup> without dietary cholesterol (Group 2) produced about the same degree of plaque as 10%  $\beta$ -casein A<sup>1</sup> or A<sup>2</sup> with dietary cholesterol (Groups 5 and 6) (see Fig. 2A). When the thoracic and abdominal segments of aorta were analysed separately, the same overall results were obtained.

When histological sections of the aortic arch were analysed morphometrically, there was no measurable intimal thickening in the presence of 20% whey alone (Group 1) or 10%  $\beta$ -casein A<sup>2</sup> with no dietary cholesterol (Group 3) (Fig. 2B). The intima to media ratio in rabbits fed 10%  $\beta$ -casein A<sup>1</sup> in the absence of dietary cholesterol (Group 2) was significantly higher than both of these groups, and produced almost the same degree of intimal thickening as 20% whey with 0.5% dietary cholesterol (Group 4) (0.04±0.01 vs 0.03±0.01,  $P < 0.05$ ). Of note, in the presence of dietary cholesterol, the intima to media ratio in the aortae of Group 7 rabbits (5%  $\beta$ -casein A<sup>1</sup>) was significantly higher than in Group 8 (5%  $\beta$ -casein A<sup>2</sup>) (Table 1). Also, the value in Group 9 rabbits (20%  $\beta$ -casein A<sup>1</sup>) was significantly higher than in rabbits fed 20%  $\beta$ -casein A<sup>2</sup> (Group 10). Groups 8 and 10 (5 and 20%  $\beta$ -casein A<sup>2</sup> plus dietary

cholesterol) had no measurable intimal thickening, while Group 6 (10% A<sup>2</sup>  $\beta$ -casein with dietary cholesterol) had a mean intima to media ratio of only 0.01, with four out of the six rabbits in this group having 0%. All  $\beta$ -casein A<sup>1</sup> groups had higher intima to media ratios than groups fed  $\beta$ -casein A<sup>2</sup> ( $P < 0.05$ ) and had intima to media ratios not significantly different from Group 4 (20% whey with 0.5% cholesterol).

### 3.5. Advanced lesions in carotid arteries

Balloon catheter injury to the right carotid artery at  $t = 0$  induced a hyperplastic neointimal thickening in all rabbits by 6 weeks (Fig. 2C). Measurement of intima to media ratios showed that in the absence of dietary cholesterol there was a slight, but not significant, increase in neointimal thickening in Group 2 rabbits (10%  $\beta$ -casein A<sup>1</sup>) compared with Group 3 (10%  $\beta$ -casein A<sup>2</sup>) and Group 1 (20% whey). Likewise, the groups fed 5, 10 and 20%  $\beta$ -casein A<sup>1</sup> plus 0.5% cholesterol had thicker neointimas than Group 1. Groups fed 5, 10 and 20%  $\beta$ -casein A<sup>2</sup> plus 0.5% cholesterol had thinner neointimas than their  $\beta$ -casein A<sup>1</sup> counterparts (see Fig. 2C, Table 1).

There was no neointimal thickening observed in any of the contralateral uninjured carotid arteries.

## 4. Discussion

When rabbits are fed a cholesterol-enriched diet, subendothelial accumulations of fat-filled macrophages (“foam cells”) are developed in most arteries [22]. These lesions are termed fatty streaks and closely resemble human juvenile fatty streaks that are present in early childhood and are considered the precursors of advanced atherosclerotic plaques [23]. Since juvenile fatty streaks are first seen in infants under 3 years of age [23], the question was raised: is casein, in addition to milk fat, responsible for fatty streak formation? Casein (as an unspecified mixture of variants) has been shown to




Fig. 2. (A) Percent luminal surface area of aorta covered by lipid-filled lesions. Note that in Group 1 there were no areas of lipophilic staining. (B) Thickness of lesions in aortic arch expressed as intima to media ratio (I:M). Note that there was no intimal thickening in Groups 1 and 3. (C) Thickness of lesions in balloon injured carotid artery expressed as I:M. Group 1: 20% whey diet; Group 2: 10%  $\beta$ -casein A<sup>1</sup>+10% whey; Group 3: 10%  $\beta$ -casein A<sup>2</sup>+10% whey; Group 4: 20% whey+0.5% cholesterol; Group 5: 10%  $\beta$ -casein A<sup>1</sup>+0.5% cholesterol; and Group 6: 10%  $\beta$ -casein A<sup>2</sup>+0.5% cholesterol. \* $P < 0.05$ .

promote atherosclerosis in several animal models, including rabbits [16,24], monkeys [17] and mice [18]. None of these studies, however, defined which particular casein variant was responsible.

In the present study, using a rabbit model it was found that both  $\beta$ -casein A<sup>1</sup> and A<sup>2</sup> led to aortic fatty streaks but A<sup>1</sup> produced considerably more extensive lesions than both A<sup>2</sup> and whey. Indeed, 10% A<sup>1</sup> without dietary cholesterol produced about the same luminal surface area of the aorta covered by lesions as 10% A<sup>2</sup>

with added dietary cholesterol. The thickness of fatty streaks in the aortic arch of rabbits fed A<sup>1</sup> was significantly greater than that in rabbits fed A<sup>2</sup> or whey. Even in the presence of 0.5% dietary cholesterol, the thickness of these lesions was significantly greater in all A<sup>1</sup> than A<sup>2</sup> groups. Thus A<sup>1</sup> is more atherogenic than both the whey and the A<sup>2</sup> casein variant.

A secondary aspect of this study was to investigate the potential of  $\beta$ -casein A<sup>2</sup> to exert an athero-protective effect compared with whey in the presence of 0.5%

dietary cholesterol. Only 20% A<sup>2</sup> produced a smaller surface area of aorta covered by fatty streaks than 20% whey under these conditions. However, aortic arch fatty streak thickness was significantly smaller for both 20 and 5% A<sup>2</sup> fed animals compared with 20% whey. These results, combined with the fact that 10% A<sup>2</sup> with no added cholesterol produced a significantly lower serum cholesterol level and LDL to HDL ratio than all other groups including whey only and 10% A<sup>1</sup>, demonstrate for the first time that  $\beta$ -casein A<sup>2</sup> has a mildly athero-protective effect while  $\beta$ -casein A<sup>1</sup> is most definitely atherogenic.

These results are consistent with epidemiological studies that suggest a strong relationship between mortality from cardiovascular disease and consumption of  $\beta$ -casein A<sup>1</sup>, even though other studies show no association between reported milk consumption per se and cardiovascular mortality [11]. In the states of the former West Germany, where allelic composition of regional cattle herds has remained virtually constant since 1950s, IHD mortality by state correlates with the estimated consumption of the  $\beta$ -casein A<sup>1</sup> allele [12]. Also, the IHD mortality rates of the populations of Toulouse in France and Belfast in northern Ireland vary more than threefold, despite having nearly identical traditional risk factors for heart disease. Interestingly, people from Belfast are estimated to consume 3.23 times more  $\beta$ -casein A<sup>1</sup> than those from Toulouse. The communities recorded as being essentially free from IHD but also drink milk, the Masai [13] and Samburu [14] of Africa, obtain their milk from Zebu cattle, which produce only the  $\beta$ -casein A<sup>2</sup> allele and no  $\beta$ -casein A<sup>1</sup> [15].

What are the possible mechanisms by which the different casein variants affect the development of atherosclerosis? Firstly, there is the effect of  $\beta$ -casein A<sup>2</sup> on lowering serum cholesterol. Secondly,  $\beta$ -casein A<sup>1</sup>, but not  $\beta$ -casein A<sup>2</sup>, releases  $\beta$ -casomorphin-7 ( $\beta$ -CM-7) [25], which is thought to be involved in the oxidation of LDL through a peroxidase-dependent process.  $\beta$ -Casomorphins have been detected in the plasma of newborn calves [26] and in the small intestines of humans after cow milk consumption [27]. Pancreatic elastase releases the carboxyl terminus of  $\beta$ -CM-7 by degrading the peptide bond between Ile<sup>66</sup> and His<sup>67</sup> of  $\beta$ -casein A<sup>1</sup>. However, elastase is unable to cleave the bond between Ile<sup>66</sup> and Pro<sup>67</sup> of  $\beta$ -casein A<sup>2</sup>, and so  $\beta$ -CM-7 cannot be released. Analysis of protein oxidation products isolated in atherosclerotic lesions has implicated the tyrosyl radical (among others) in LDL oxidation [28]. Torreilles and Guerin [29] found that bovine casein-derived peptides with tyrosyl-end residues (such as  $\beta$ -CM-7) could promote peroxidase-dependent oxidation of human LDL, implying that the tyrosyl ending peptide is a diffusible catalyst that conveys oxidising potential to LDL lipids. Finally, the fact that

there was no difference in intima to media ratio in the balloon injured right carotid artery of rabbits fed A<sup>1</sup>, A<sup>2</sup> or whey may point to possible mechanistic differences. Endothelium was present in the aorta but absent in the right carotid artery, and this suggests different effects on the endothelium following ingestion of the two  $\beta$ -caseins, perhaps in relation to expression of adhesion molecules such as ICAM-1 which in turn leads to monocyte/macrophage invasion of the vessel wall [30,31].

What might the differential effects of the casein variants have on the wider community?  $\beta$ -Casein allele frequency data indicate that of the four major milk-producing breeds of cow in the Western world, half produce milk that is rich in  $\beta$ -casein A<sup>1</sup> and only one of the four breeds produces milk that is almost purely A<sup>2</sup>. This means that the milk sold in supermarkets and other stores will contain a high proportion of the  $\beta$ -casein A<sup>1</sup> allele. Since children develop juvenile fatty streaks as a precursor to more advanced lesions in later life, the type of casein ingested in childhood may be critical to the likelihood of developing atherosclerosis. The public could be offered the healthier choice of pure A<sup>2</sup> milk, which would be particularly important for the children of persons with known cardiovascular disease. Since the milk from Guernsey cows contains 98% A<sup>2</sup>, it may be possible to change herd allelic composition by 'breeding out' those cows with A<sup>1</sup>/A<sup>1</sup> alleles. These measures may thus contribute to improve human health by reducing a risk factor that contributes to ischaemic heart disease, the leading cause of premature death in the developed world today.

## References

- [1] Davies DT, Law AJR. The content and composition of creamery milks in south-west Scotland. *J Dairy Res* 1980;47:83–90.
- [2] Jenness R. Comparative aspects of milk proteins. *J Dairy Res* 1979;46:197–210.
- [3] Whitney RMCL, Brunner JR, Ebner KE, Farrell HM, Jr., Josephson RV, Morr CV, Swaisgood HE. Nomenclature of the protein of cow's milk: fourth revision. *J Dairy Sci* 1976;59:795–815.
- [4] Stewart AF, Bonsing J, Beattie CW, Shah F, Willis IM, Mackinlay AG. Complete nucleotide sequences of bovine  $\alpha_{s2}$ - and  $\beta$ -casein cDNAs: comparisons with related sequences in other species. *Mol Biol Evol* 1987;4:231–41.
- [5] Formaggioni P, Summer A, Malacarne M, Mariani P. Milk protein polymorphism: detection and diffusion of the genetic variants in *Bos* genus, Vol. XIX. Università degli Studi di Parma, Annali della Facoltà di Medicina Veterinaria, 1999.
- [6] Li FH, Gaunt SN. A study of genetic polymorphisms of milk beta-lactoglobulin, alpha s1-casein, beta-casein and kappa-casein in five dairy breeds. *Biochem Genet* 1972;6:9–20.
- [7] Barroso A, Dunner S, Canon J. Technical note: use of PCR-single-strand conformation polymorphism analysis for detection of bovine  $\beta$ -casein variants A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and B. *J Anim Sci* 1999;77:2629–32.

- [8] Pinder SJ, Perry BN, Skidmore CJ, Savva D. Analysis of polymorphism in the bovine casein genes by use of the polymerase chain reaction. *Anim Genet* 1991;22:11–20.
- [9] McLean DM, Graham ERB, Ponzoni RW, McKenzie HA. Effects of milk protein genetic variants on milk yield and composition. *J Dairy Sci* 1984;51:531–46.
- [10] Baev AA, Smirnov IK, Gorodetskii SI. Primary structure of bovine  $\beta$ -casein cDNA. *Mol Biol (NY)* 1987;21:214–22.
- [11] Ness AR, Davey Smith G, Hart C. Milk, coronary heart disease and mortality. *J Epidemiol Commun Health* 2001;55:379–82.
- [12] McLachlan CNS.  $\beta$ -Casein A<sup>1</sup>, ischaemic heart disease mortality and other illnesses. *Med Hypotheses* 2001;56:262–72.
- [13] Biss K, Ho K-J, Mikkelsen BS, Lewis L, Taylor CB. Some unique biological characteristics of the Masai of East Africa. *N Engl J Med* 1971;28:694–9.
- [14] Shaper AG. Cardiovascular studies in the Samburu tribe of northern Kenya. *Am Heart J* 1962;63:437–42.
- [15] Aschaffenburg R, Sen A, Thompson MP. Genetic variants of casein in Indian and African Zebu cattle. *Comp Biochem Physiol* 1968;25:177–84.
- [16] Kratky RG, Ivey J, Rogers KA, Daley S, Roach MR. The distribution of fibro-fatty atherosclerotic lesions in the aortae of casein- and cholesterol-fed rabbits. *Atherosclerosis* 1993;99:121–31.
- [17] Anthony MS, Clarkson TB, Williams JK. Effects of soy isoflavones on atherosclerosis: potential mechanisms. *Am J Clin Nutr* 1998;68:1390S–3S.
- [18] Ni W, Tsuda Y, Sakono M, Imaizumi K. Dietary soy protein isolate, compared with casein, reduces atherosclerotic lesion area in apolipoprotein E-deficient mice. *J Nutr* 1998;128:1884–9.
- [19] Beynen AC, Van de Meer R, West CE. Mechanisms of casein-induced hypercholesterolemia: primary and secondary features. *Atherosclerosis* 1986;60:291–3.
- [20] Park MS, Kudchodkar BJ, Leipa GU. Effects of dietary animal and plant proteins on cholesterol metabolism in immature and mature rats. *J Nutr* 1987;117:30–5.
- [21] Manderson JA, Mosse PRL, Safstrom JA, Young SB, Campbell GR. Balloon catheter injury to rabbit carotid artery. I. Changes in smooth muscle phenotype. *Arteriosclerosis* 1989;9:289–98.
- [22] Walker LN, Reidy MA, Bowyer DE. Morphology and cell kinetics of fatty streak lesion formation in the hypercholesterolemic rabbit. *Am J Pathol* 1986;125:450–9.
- [23] Wissler RW, Robertson AL, Cornhill JF, McGill HC, McMahan CA, Strong JP. Natural history of aortic and coronary atherosclerotic lesions in youth: findings from the PDAY study. *Arterioscler Thromb* 1993;13:1291–8.
- [24] Katan MB, Vroomen LHM, Hermus RJJ. Reduction of casein-induced hypercholesterolemia and atherosclerosis in rabbits and rats by dietary glycine, arginine and alanine. *Atherosclerosis* 1982;43:381–91.
- [25] Jinsmaa Y, Yoshikawa M. Enzymatic release of neocasomorphin and  $\beta$ -casomorphin from bovine  $\beta$ -casein. *Peptides* 1999;20:957–62.
- [26] Svedberg J, De Haas J, Leimenstoll KG, Paul F, Teschemacher H. Demonstration of  $\beta$ -casomorphin immunoreactive material in *in vitro* digests of bovine milk and in small intestine contents after bovine milk ingestion in adult humans. *Peptides* 1985;6:825–30.
- [27] Meisel H. Chemical characterisation and opioid activity of an exorphin isolated from *in vivo* digests of casein. *FEBS Lett* 1986;196:223–7.
- [28] Heinecke JW. Mass spectrometric quantification of amino acid oxidation products in proteins: insights into pathways that promote LDL oxidation in the human artery wall. *FASEB J* 1999;13:1113–20.
- [29] Torrealles J, Guerin M-C. Casein-derived peptides can promote human LDL oxidation by a peroxidase-dependent and metal-independent process. *French CR Soc Biol* 1995;189:933–42.
- [30] Poston RN, Haskard DO, Coucher JR, Gall NP, Johnson-Tidey RR. Expression of intercellular adhesion molecule-1 in atherosclerotic plaques. *Am J Pathol* 1992;140:665–73.
- [31] Van der Wal AC, Das PK, Tigges AJ, Becker AE. Adhesion molecules on the endothelium and mononuclear cells in human atherosclerotic lesions. *Am J Pathol* 1992;141:1427–33.